

REPORT DOCUMENTATION PAGEForm Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. **PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**

1. REPORT DATE (DD-MM-YYYY)

03-24-2004

2. REPORT TYPE

Technical Paper (View Graph)

3. DATES COVERED (From - To)**4. TITLE AND SUBTITLE**

Polynitrogen Chemistry: Recent Development in Pentazole and Polyazide Chemistry

5a. CONTRACT NUMBER

F04611-99-C-0025

5b. GRANT NUMBER**5c. PROGRAM ELEMENT NUMBER****6. AUTHOR(S)**

Ashwani Vij, Vandan Vij, R. Haiges, Karl O. Christe

5d. PROJECT NUMBER

DARP

5e. TASK NUMBER

A205

5f. WORK UNIT NUMBER**7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)**ERC Incorporated
555 Sparkman Drive
Huntsville, AL 35816-0000**8. PERFORMING ORGANIZATION REPORT NUMBER****9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)**Air Force Research Laboratory (AFMC)
AFRL/PRSB
4 Draco Drive
Edwards AFB CA 93524-7160**10. SPONSOR/MONITOR'S ACRONYM(S)****11. SPONSOR/MONITOR'S NUMBER(S)**

AFRL-PR-ED-VG-2004-088

12. DISTRIBUTION / AVAILABILITY STATEMENT

Approved for public release; distribution unlimited.

13. SUPPLEMENTARY NOTESAmerican Chemical Society Meeting (ACS) National Meeting
Anaheim, CA 28 Mar - 01 Apr 2004**14. ABSTRACT**

20040503 192

15. SUBJECT TERMS**16. SECURITY CLASSIFICATION OF:****17. LIMITATION OF ABSTRACT****18. NUMBER OF PAGES****19a. NAME OF RESPONSIBLE PERSON**
Linda Talon**a. REPORT****b. ABSTRACT****c. THIS PAGE**

Unclassified

Unclassified

Unclassified

A

38

19b. TELEPHONE NUMBER (include area code)
(661) 275-5865Standard Form 298 (Rev. 8-98)
Prescribed by ANSI Std. Z39.18

Polynitrogen Chemistry: Recent Development in Pentazole and Polyazide Chemistry



Ashwani Vij


Space and Missile Propulsion Division
Air Force Research Laboratory/PRSP
Edwards AFB, CA 93524
ashwani.vij@edwards.af.mil
(661) 275-6278

Inorganic Division, 227th National ACS Meeting, Anaheim-CA, April 1, 2004
Approved for public release, distribution unlimited



Why Polynitrogen Compounds ?




- Polynitrogen compounds contain only nitrogen atoms and are expected to have unusual properties. Most important among these are:
 - *High endothermicity*
 - *"Green" propellant*
"combustion" product is only gaseous N₂
 - *High density*
 - *High I_{sp} values when compared to other monopropellants or bipropellants*
 -  *High detonation velocity*




Geometry of the N_5^+ cation



V-Shaped Geometry




Calculated Structure




Experimental Structure

Vij, Wilson, Vij, Tham, Sheehy & Christie,
J. Am. Chem. Soc., 2001, 123, 6308-6313



Resonance Structure



N2 makes contacts at 2.723 and 2.768 Å
N4 contacts are at 2.887 and 2.814 Å

C&E News, 2000, 78, 41

April 1, 2004

Inorganic Division – Main Group Chemistry, 227th National ACS Meeting, Anaheim-CA, April 1, 2004
Approved for public release, distribution unlimited




3



(In)Compatability of N_5^+



Attempts to couple N_5^+ with energetic anions
can result in explosive reactions !!!



Our goal is the synthesis of an “aromatic” polynitrogen anion with

- A high first ionization potential
- A high activation energy barrier towards decomposition


April 1, 2004

Inorganic Division – Main Group Chemistry, 227th National ACS Meeting, Anaheim-CA, April 1, 2004
Approved for public release, distribution unlimited

4




New Polynitrogen Anions as Counterparts for N_5^+



Pentazole anion (N_5^-)

- Theoretical calculations show that this anion has a 28 kcal/mole activation energy barrier for decomposition and its decomposition to N_3^- and N_2 is only 11 kcal/mol exothermic
- Free pentazole has not been isolated to date. Only aryl substituted pentazoles can be isolated and stabilized at low temperatures. These compounds rapidly decompose above 273K to form aryl azides and N_2 gas



April 1, 2004

Inorganic Division – Main Group Chemistry, 227th National ACS Meeting, Anaheim-CA, April 1, 2004
Approved for public release, distribution unlimited

5




Synthetic Challenge – How do we make These New Anions??



Synthesis of Substituted Pentazoles

Sources for the Pentazole Anion (N_5^-)



April 1, 2004

Inorganic Division – Main Group Chemistry, 227th National ACS Meeting, Anaheim-CA, April 1, 2004
Approved for public release, distribution unlimited


6




Formation and Stability of Silyl Diazonium Salts



- Failed attempts to synthesize silyl diazonium salts



- R_3SiN_2^+ salts are unstable and spontaneously lose N_2



Theoretical calculations support this experimental observation.

April 1, 2004

Inorganic Division – Main Group Chemistry, 227th National ACS Meeting, Anaheim-CA, April 1, 2004
Approved for public release, distribution unlimited


7



Synthesis of Aryldiazonium Salts




Aqueous Media



Non-aqueous Media

$\text{R} = \text{H, OH, OCH}_3, \text{OC}_6\text{H}_5, \text{OC}_6\text{H}_4\text{N}_2^+, \text{N}(\text{CH}_3)_2$






April 1, 2004

Inorganic Division – Main Group Chemistry, 227th National ACS Meeting, Anaheim-CA, April 1, 2004
Approved for public release, distribution unlimited

8



Pentazole Formation... Not a Trivial Chore !!!



April 1, 2004

Inorganic Division – Main Group Chemistry, 227th National ACS Meeting, Anaheim-CA, April 1, 2004
Approved for public release, distribution unlimited

9




Identification of Arylpentazoles



Pentazoles can be characterized by low temperature NMR spectral studies using ^{15}N labeled samples.

- ^1H NMR: AB-type spectrum with H_a and H_b at 8.0 and 7.0 ppm
- ^{14}N NMR: N_1 at ~ -80 ppm
- ^{15}N NMR: N_2/N_5 at ~ -27 ppm and N_3/N_4 at ~ -4 ppm



Note: Qualitative evidence for the presence of a pentazole ring: N_2 gas evolution in solution

April 1, 2004

Inorganic Division – Main Group Chemistry, 227th National ACS Meeting, Anaheim-CA, April 1, 2004
Approved for public release, distribution unlimited

10



Cleavage of the Aryl-Pentazole Bond with Retention of the Pentazole Ring



Chemical Methods


- Ozonolysis does not work! (Ugi, Radziszewski)

V. Benin, P. Kszynski and G. J. Radziszewski, *J. Org. Chem.*, 2002, 67, 1354

- Nucleophilic substitution using strong nucleophiles such as the OH^- , OR^- , F^- etc.

Collisional Fragmentation (ElectroSpray Ion Mass Spectroscopy – ESIMS)

- Electrospray is very gentle and produces high concentration of the parent anion which can be mass selected
- Negative ion detection eliminates interference from neutral or positively charged species




April 1, 2004

Inorganic Division – Main Group Chemistry, 227th National ACS Meeting, Anaheim-CA, April 1, 2004
Approved for public release, distribution unlimited

11



MSMS of the Parent Ion Peak



Vij et al., *Angew. Chem.* 2002, 41, 3051

m/z

G&E News, 2002, 80, 8


April 1, 2004

Inorganic Division – Main Group Chemistry, 227th National ACS Meeting, Anaheim-CA, April 1, 2004
Approved for public release, distribution unlimited

12



Chemical Cleavage of the C-N Bond




April 1, 2004

Inorganic Division - Main Group Chemistry, 227th National ACS Meeting, Anaheim-CA, April 1, 2004
Approved for public release, distribution unlimited

13



Aryl-pentazole bond cleavage: N_5^- anion in Solution




April 1, 2004

Inorganic Division - Main Group Chemistry, 227th National ACS Meeting, Anaheim-CA, April 1, 2004
Approved for public release, distribution unlimited

14



Subsequent Decomposition of the Pentazole Anion



Current Data Parameters
NAME Pyrazacore
EXPNO 25
PROCNO 1

F2 - Acquisition Parameters
Date_ 20030709
Time 12.27
INSTRUM spect
PROBHD 5 mm BBO 1H-
PULPROG zgpg30
TD 32768
SOLVENT CDCl3
NS 0
DS 0
SWH 20325.203 Hz
FIDRES 0.430274 Hz
AQ 0.8061428 sec
RG 57344
DW 24.400 usec
DE 6.00 usec
TE 300.0 K
D1 60.0000000 sec

***** CHANNEL f1 *****
NUC1 15N
P1 12.50 usec
PL1 -4.00 dB
SFO1 40.543250 MHz

F2 - Processing parameters
SI 16384
SF 40.5407913 MHz
WDW EM
SSB 0
LB 0.25 Hz
GB 0
PC 1.00

April 1, 2004

Inorganic Division - Main Group Chemistry, 227th National ACS Meeting, Anaheim-CA, April 1, 2004
Approved for public release, distribution unlimited


15



Pentazoles with Heterocyclic Substituents



- Tetrazolyl system is unstable above -70 °C and the pentazole ring rapidly decomposes to liberate N₂ gas.



A. Hammerl and T. M. Klapoetke, *Inorg. Chem.* 2002, 41, 906-912

- In comparison, the pentazole ring derived from 2-amino-4,5-dicyanoimidazole shows higher thermal stability (-30 °C)




April 1, 2004

Inorganic Division - Main Group Chemistry, 227th National ACS Meeting, Anaheim-CA, April 1, 2004
Approved for public release, distribution unlimited

16



¹⁵N NMR of 2-pentazoly-4,5-dicyanoimidazole



¹⁵N NMR recorded in a mixture of methanol and acetonitrile at -30 °C, nitromethane used as an external reference (0 ppm)


April 1, 2004

Inorganic Division – Main Group Chemistry, 227th National ACS Meeting, Anaheim-CA, April 1, 2004
Approved for public release, distribution unlimited

17



Pentazolate Anion in Solution ?



- ¹⁵N NMR shows a peak at ~ -10 ppm (-30 °C) upon addition of base, which slowly decomposes to form N₂ and azide ion.
- This peak is also observed upon adding a base to the solution of arylpentazoles at -30 °C.

April 1, 2004

Inorganic Division – Main Group Chemistry, 227th National ACS Meeting, Anaheim-CA, April 1, 2004
Approved for public release, distribution unlimited

18



Chronology of the Pentazolate Anion



- ESIMS of *para*-hydroxyphenylpentazole
Vij, Pavlovich, Wilson, Vij, Christie, Angew. Chem. Intl. Ed. Engl 2002, 41, 3051
Submitted: April 30, 2002; accepted July 3, 2002
- ^{15}N NMR studies showing a peak at -10.2 ppm (-40°C) due to the Pentazole anion resulting from cleavage of *para*-methoxyphenylpentazole which slowly decomposes to form N_2 and azide ion. Upon standing for several days, all peaks disappear!
Butler, Stephens & Burke, Chem. Commun. 2003, 1016
Submitted: February 6, 2003; accepted February 27, 2003
- Laser Desorption Ionization (LDI) time-of-flight (TOF) mass spectrometry of solid *para*- N,N -dimethylaminophenylpentazole shows peaks at m/z : 70 (N_5^-) and -42 (N_3^-). Peak at 70 confirmed by ^{15}N labeling experiment.
Ostmark, Wallin, Brinck, Carlqvist, Claridge, Hedlund & Yudina, Chem. Phys. Lett., 2003, 379, 539
Submitted: Jun. 27, 2003; accepted August 27, 2003


April 1, 2004

Inorganic Division – Main Group Chemistry, 227th National ACS Meeting, Anaheim-CA, April 1, 2004
Approved for public release, distribution unlimited

19



What are “normal” N-N distances in azides?



April 1, 2004

Inorganic Division – Main Group Chemistry, 227th National ACS Meeting, Anaheim-CA, April 1, 2004
Approved for public release, distribution unlimited

20



Abnormalities in azide distances: Artifacts or Structural Contaminants?



- ✓ A large number of crystal structures reported in Cambridge/Inorganic CSD report unusually short $N_{\alpha}-N_{\beta}$ (0.8-1.0 Å) and long $N_{\beta}-N_{\gamma}$ (1.2-1.4) distances.

According to VB theory, in covalently bonded azides,
 $N_{\alpha}-N_{\beta} > N_{\beta}-N_{\gamma}$

Wolfgang, F. and Klapoetke, T. In *Inorganic Chemistry Highlights*; Meyer, G., Naumann, D. and Wesemann, L. Eds.; Wiley-VCH: Weinheim, 2002, Chapter 16 and references therein

- ✓ In most cases, these derivatives were prepared from metal chloride salts and/or recrystallized from chlorinated solvents


April 1, 2004

Inorganic Division – Main Group Chemistry, 227th National ACS Meeting, Anaheim-CA, April 1, 2004
Approved for public release, distribution unlimited

21



Reactivity of hexachloroantimonate (VI) with Trimethylsilylazide



- ✓ The substitution of all the six chlorine atoms in SbCl_6^- by the azide groups could not be accomplished in a single step, as reported in literature. The stepwise substitution gives a good insight into the substitution mechanism.

- ✓ Total substitution was achieved after four “refreshment” cycles of the reagents. During the intermediate cycles, the azide content gradually increased from two to five.


April 1, 2004

Inorganic Division – Main Group Chemistry, 227th National ACS Meeting, Anaheim-CA, April 1, 2004
Approved for public release, distribution unlimited


22



The tetraphenylarsonium antimonychloride azide case



Orthorhombic, $P2_12_12_1$
 a, b, c (Å) = 7.7744(14), 13.610(3), 27.094(5)
 $V = 2866.8(10)$, $Z = 4$
 $R = 3.22\%$, $S = 1.088$, highest peak = $0.58 \text{ e}^-/\text{\AA}^3$
Flack's parameter = 0.03(1)



Chloride contamination (%)
NI-N3 = 10; N4-N6 = 18; N7-N9 = 16; NI0-N12 = 37
 $R = 3.22\%$, $S = 1.088$, highest peak = $0.58 \text{ e}^-/\text{\AA}^3$
Flack's parameter = 0.03(1)

April 1, 2004

Inorganic Division - Main Group Chemistry, 227th National ACS Meeting, Anaheim-CA, April 1, 2004
Approved for public release, distribution unlimited

23



Summary



- Synthesized aryl pentazoles: hydroxy group at the *para*-position on the aryl ring gives the best results as observed during this study.
- Demonstrated selective cleavage of C-N bond by ESIMS with retention of pentazole ring. Results confirmed studying ^{15}N labeled pentazoles.
- Experimental detection of pentazolate anion in solution using different substrates.
- Offers potential pathway for bulk synthesis of N_5^- salts
- Chloride ion cause abnormalities in N-N bonds in azides


April 1, 2004

Inorganic Division - Main Group Chemistry, 227th National ACS Meeting, Anaheim-CA, April 1, 2004
Approved for public release, distribution unlimited

24



Acknowledgments



Dr. Karl Christe, Dr. William Wilson, Ms. Vandana Vij (ERC Inc.)

Dr. Gregory Drake (AFRL)



Dr. Ralf Haiges (USC)

Dr. Fook Tham (UC Riverside)



Dr. James Pavlovich (UCSB)

Dr. Robert Corley, Dr. Ronald Channell, Mr. Michael Huggins (AFRL)



\$\$\$

Dr. Don Woodbury, Dr. Arthur Morrish (DARPA)

Dr. Michael Berman (AFOSR)

April 1, 2004

Inorganic Division - Main Group Chemistry, 227th National ACS Meeting, Anaheim-CA, April 1, 2004
Approved for public release, distribution unlimited

25



BACKUP

April 1, 2004

Inorganic Division - Main Group Chemistry, 227th National ACS Meeting, Anaheim-CA, April 1, 2004
Approved for public release, distribution unlimited

26




Recipe for Synthesizing Neutral Polynitrogen Compounds



- Combine a polynitrogen cation with a polynitrogen anion to form a neutral polynitrogen compound.




ONLY TWO STABLE POLYNITROGEN IONS KNOWN TO EXIST IN BULK



N_5^+ cation

(discovered in 1999, AFRL, Christie)



N_3^- anion

(discovered in 1890, Curtius)



April 1, 2004

Inorganic Division – Main Group Chemistry, 227th National ACS Meeting, Anaheim-CA, April 1, 2004
Approved for public release, distribution unlimited

33



Episode I...Generation of the starting material




April 1, 2004

Inorganic Division – Main Group Chemistry, 227th National ACS Meeting, Anaheim-CA, April 1, 2004
Approved for public release, distribution unlimited


34




Episode II....cis- or trans- disubstitution with azide groups?



cis-isomer



trans-isomer




April 1, 2004

Inorganic Division – Main Group Chemistry, 227th National ACS Meeting, Anaheim-CA, April 1, 2004
Approved for public release, distribution unlimited


35




Episode III...Substitution of 3rd chlorine... fac- or mer- isomer ???



mer-SbCl₃(N₃)₃



fac-SbCl₃(N₃)₃



April 1, 2004

Inorganic Division – Main Group Chemistry, 227th National ACS Meeting, Anaheim-CA, April 1, 2004
Approved for public release, distribution unlimited


36



Episode VI... Complete substitution of chlorine atoms



No crystal structure obtained yet. However, IR and Raman spectroscopy shows that Sb-Cl bonds are absent i.e., complete substitution by the azide groups.



April 1, 2004

Inorganic Division – Main Group Chemistry, 227th National ACS Meeting, Anaheim-CA, April 1, 2004
Approved for public release, distribution unlimited


27



ESIMS of para-Phenoxypentazole



Observed peaks in the MSMS of 162



Vij, Pavlovich, Wilson, Vj & Christe, *Angew. Chem. Int. Ed.*, 2002, 41, 3051-3054


April 1, 2004

Inorganic Division – Main Group Chemistry, 227th National ACS Meeting, Anaheim-CA, April 1, 2004
Approved for public release, distribution unlimited

28



Episode V: Synthesis of Chloropentaazidoantimonate(VI) Anion



The Structure of $\text{Ph}_4\text{PSbCl}(\text{N}_3)_5$

- ✓ The crystals grown from CH_3CN
- ✓ Triclinic space group $P-1$
- ✓ Cell constants: $a = 11.134(3) \text{ \AA}$, $b = 11.663(3) \text{ \AA}$, $c = 13.754(4) \text{ \AA}$; $\alpha = 104.314(5)^\circ$, $\beta = 97.914(5)^\circ$, $\gamma = 115.807(4)^\circ$
- ✓ $Z = 2$
- ✓ $R = 0.0762$
- ✓ All azide distances "normal" except N10-N11-N12


April 1, 2004

J. 227th National ACS Meeting, Anaheim-CA, April 1, 2004
Approved for public release, distribution unlimited

37



Reaction with Trimethylsilyl Azide



$\text{X} = \text{N}, \text{O}$

No pentazoles were isolated !!!

Reactions carried out in acetonitrile at -30°C

April 1, 2004

Inorganic Division – Main Group Chemistry, 227th National ACS Meeting, Anaheim-CA, April 1, 2004
Approved for public release, distribution unlimited

38